

The authors and reviewers have made every attempt to ensure the information in the Family Medicine Clinical Cards is correct - it is possible that errors may exist. Accordingly, the source references or other authorities should be consulted to aid in determining the assessment and management plan of patients. The Cards are not meant to replace customized patient assessment nor clinical judgment. They are meant to highlight key considerations in particular clinical scenarios, largely informed by relevant guidelines in effect at the time of publication. The authors cannot assume any liability for patient outcomes when these cards are used.

Canadian Family Medicine Clinical Card

A16 2011
www.learnfm.ca

Bach TV
O'Beirne M
Keegan DA

Common Prenatal Problems

NAUSEA AND VOMITING

- begins @ 6 wks, peaks @ 9 wks; 60% resolve by 12 wks, 91% by 20 wks, 5% entire preg
- women with N&V have fewer spont. abortions and stillbirths vs. women without N&V
- hyperemesis gravidarum = most severe form of NV occurs in < 1%

1st line treatment

Start Diclectin (combo of 10 mg doxylamine + 10 mg pyridoxine)

- recommended dose = 4 tabs daily (2 qhs + 1 qam + 1 qafternoon)
- up to 8 tabs daily, adjust prn, delayed action (takes 8 hr to work)

2nd line treatment

Add or switch to a substitute: antihistamines, e.g. dimenhydrinate, diphenhydramine

- for acute or breakthrough NV, use IV and PR formulation

3rd line treatment

If **dehydrated**:

- **warning signs: wt loss, oliguria**
- hospitalize with IV fluid replacement, multivitamin IV, antiemetic IV

If **well-hydrated**, add or switch to a substitute (in order of fetal safety):

- phenothiazines, e.g. chlorpromazine; metoclopramide; ondansetron

4th line treatment

Corticosteroids, e.g. methylprednisolone, consider only in refractory cases

- avoid corticosteroids at \leq 10 wks because of higher risk of oral clefting

Consider other causes or exacerbating factors, test:

- electrolytes, Cr, Bun, liver function, TSH, drug levels, U/S and *H. pylori* testing

Notes

Diet and lifestyle Δ s, including:

- eat what appeals, avoid triggers, smaller frequent meals, rest plenty
- stop prenatal multivitamin with Fe (Fe causes gastric irritation/ N&V)

Adjuvant treatment can be added at any time, including:

- ginger supp (in any form, maximum dose = < 1 g per day)
- pyridoxine, acupressure, acupuncture

HEARTBURN AND ACID REFLUX

1st line Antacids (avoid Mg trisilicate and bicarbonate-containing antacids)

2nd line - H2 antagonists, e.g. ranitidine

- PPIs, e.g. omeprazole, pantoprazole

AVOID Pepto Bismol because of salicylate absorption

Notes Lifestyle modifications, including: eat smaller and more frequent meals, avoid eating near bedtime, elevate head of bed

URINARY TRACT INFECTION

- treat asympt. bacteriuria; if not, \uparrow risk of cystitis, pyelonephritis & preterm labour

1st line Penicillins, cephalosporins, fluoroquinolones, nitrofurantoin, phenazopyridine

AVOID - nitrofurantoin \geq 38 wks \rightarrow hemolytic anemia in fetus or newborn
- TMP-SMX in first trimester \rightarrow neural tube defects
- TMP-SMX \geq 32 wks \rightarrow increased kernicterus in newborn
- tetracycline / doxycycline \rightarrow deposition on bones and teeth

Notes Prophylactic treatment (if desired): vit C 500 mg daily, cranberry juice

HEADACHE

- **warning signs of severe preeclampsia: sudden onset in 3rd trimester with vision changes, RUQ pain, facial edema +/- \uparrow BP**
- treatment: increase sleep & fluid intake, acetaminophen
- **avoid NSAIDS \rightarrow teratogenic** < 12 wks, \downarrow amniotic fluid \geq 12 wks

LOW BACK PAIN

treatment:

- back exercises
- chiropractic
- physiotherapy

Key References: Arsenault M, and Lane CA. The Management of Nausea and Vomiting in Pregnancy. *SOGC Clinical Practice Guidelines Number 102*. 2002; Law R, Maltepe C, Bozzo P, Einarson A. Treatment of Heartburn and Acid Reflux Associated with Nausea and Vomiting During Pregnancy. *Can Fam Physician*. 2010;56(2):143-4. Lee M, Bozzo P, Einarson A, Karen G. Urinary Tract Infections in Pregnancy. *Can Fam Physician*. 2008;54(6):853-4.