

The authors and reviewers have made every attempt to ensure the information in the Family Medicine Clinical Cards is correct - it is possible that errors may exist. Accordingly, the source references or other authorities should be consulted to aid in determining the assessment and management plan of patients. The Cards are not meant to replace customized patient assessment nor clinical judgment. They are meant to highlight key considerations in particular clinical scenarios, largely informed by relevant guidelines in effect at the time of publication. The authors cannot assume any liability for patient outcomes when these cards are used.

Canadian Family Medicine Clinical Card

A10 2019
www.learnfm.ca

Burles K
Vaughan SD
Keegan DA



Sore Throat

APPROACH

- Viral etiology is most common (causing 80-90% of infectious pharyngitis in adults and 50-70% in children), particularly if multiple symptoms (e.g. cough, conjunctivitis, rhinorrhea, hoarseness, fever, malaise, or myalgia)
- If bacterial, Group A Streptococcus (GAS) is most common etiology → perform rapid antigen detection testing (RADT) or throat swab for GAS culture if patient scores at least 2 with the following criteria (1 point each):
 - Tonsil swelling or exudate
 - Tender/swollen anterior cervical lymph nodes
 - Fever (>38°C)
 - Cough absent
 - Age 3-14 years (if age ≥45, subtract 1 point)

⚠ RED FLAGS	Possible Diagnoses (must be ruled out)
Drizzling	epiglottitis, retropharyngeal or peritonsillar abscess
Suspicion of foreign body	foreign body
Muffled “hot potato” voice	epiglottitis
Acutely unwell/toxic	epiglottitis, retropharyngeal abscess, Diphtheria, sublingual abscess (Ludwig’s angina), infectious thrombophlebitis in the internal jugular vein (Lemierre’s syndrome)
Throat pain out of proportion to findings	epiglottitis, peritonsillar abscess
Unilaterally enlarged tonsil or uvular deviation	peritonsillar abscess
Unvacc. with thick grey/ white membrane on back of throat	Diphtheria
Oral lesions	Coxsackie virus (hand, foot, and mouth disease), Herpes, PFAPA Syndrome (periodic fever with aphthous stomatitis, pharyngitis and adenitis), Stevens-Johnson syndrome, Behcet’s syndrome, Kawasaki Disease
Adenopathy and splenomegaly	EBV (infectious mononucleosis)

TREATMENT for Infectious Pharyngitis

- Most Infectious pharyngitis can be managed with analgesia and antipyretics alone
- Delaying antibiotics until throat culture results are back is reasonable, since:
 - GAS is typically self-limited (8-10 d)
 - Delaying antibiotic treatment may prevent relapse
 - Antibiotic initiation ≤9 d after illness onset prevents Rheumatic fever (i.e. there is enough time to wait for culture results and still be effective)

Treat for GAS if: <input type="checkbox"/> +ve RADT <input type="checkbox"/> +ve throat culture	<ul style="list-style-type: none"> - Adults: Penicillin VK 600mg PO BID or 300mg PO TID x 10d - Pediatric: Penicillin VK 40mg/kg/d PO div BID x 10d - Penicillin allergy: can use Cephalexin (penicillin non-anaphylaxis), Clindamycin, Azithromycin, or Clarithromycin
--	---

If no improvement with antibiotics after 72 hrs, assess for:

- Antibiotic non-compliance
- Concurrent viral infection in GAS carrier (20% of children are carriers)
- Suppurative complications (sinusitis, retropharyngeal or peritonsillar abscess)

Key References: Shulman ST, Bisno AL, Clegg HW, et al. Clinical practice guideline for the diagnosis and management of group A streptococcal pharyngitis: 2012 update by the Infectious Diseases Society of America. *Clin Infect Dis.* 2012;55(10):1279-82. ESCMID Sore Throat Guideline Group Pelucchi C, Grigoryan L, Galeone C, et al. Guideline for the management of acute sore throat. *Clin Microbiol Infect.* 2012;Suppl 1:1-28.